

Drug Policy:

Pedmark™ (sodium thiosulfate)

POLICY NUMBER UM ONC_1467	SUBJECT Pedmark™ (sodium thiosulfate)		DEPT/PROGRAM UM Dept	PAGE 1 of 3
DATES COMMITTEE REVIEWED 11/09/22, 11/08/23	APPROVAL DATE November 8, 2023	EFFECTIVE DATE November 27, 2023	COMMITTEE APPROVAL DATES 11/09/22, 11/08/23	
PRIMARY BUSINESS OWNER: UM		COMMITTEE/BOARD APPROVAL Utilization Management Committee		
NCQA STANDARDS UM 2		ADDITIONAL AREAS OF IMPACT		
CMS REQUIREMENTS	STATE/FEDERAL REQUIREMENTS		APPLICABLE LINES OF BUSINESS Commercial, Exchange, Medicaid	

I. PURPOSE

To define and describe the accepted indications for Pedmark (sodium thiosulfate) usage in the treatment of cancer, including FDA approved indications, and off-label indications.

New Century Health (NCH) is responsible for processing all medication requests from network ordering providers. Medications not authorized by NCH may be deemed as not approvable and therefore not reimbursable.

The use of this drug must be supported by one of the following: FDA approved product labeling, CMS-approved compendia, National Comprehensive Cancer Network (NCCN), American Society of Clinical Oncology (ASCO) clinical guidelines, or peer-reviewed literature that meets the requirements of the CMS Medicare Benefit Policy Manual Chapter 15.

II. INDICATIONS FOR USE/INCLUSION CRITERIA

A. Continuation requests for a not-approvable medication shall be exempt from this NCH policy provided:

1. The requested medication was used within the last year, **AND**
2. The member has not experienced disease progression and/or no intolerance to the requested medication, **AND**
3. Additional medication(s) are not being added to the continuation request.

B. Ototoxicity Prophylaxis with Cisplatin

1. Pedmark (sodium thiosulfate) may be used to reduce the risk of ototoxicity in a pediatric member 1 month to 18 years of age and older with a localized and non-metastatic solid tumor who will be receiving cisplatin treatment **AND**
2. Pedmark (sodium thiosulfate) will be separated from cisplatin infusion by at least 6-10 hours due to the decrease in efficacy of cisplatin when both agents are used together; for this reason, Pedmark (sodium thiosulfate) may not be used with cisplatin given as a continuous infusion over 6-24 hours.

III. EXCLUSION CRITERIA

- A. Dosing exceeds single dose limit of Pedmark (sodium thiosulfate) 10 gm/m² (for weight less than 5 kg), 15 gm/m² (5-10 kg weight), or 20 gm/m² (greater than 10 kg weight).
- B. Investigational use of Pedmark (sodium thiosulfate) with an off-label indication that is not sufficient in evidence or is not generally accepted by the medical community. Sufficient evidence that is not supported by CMS recognized compendia or acceptable peer reviewed literature is defined as any of the following:
 - 1. Whether the clinical characteristics of the patient and the cancer are adequately represented in the published evidence.
 - 2. Whether the administered chemotherapy/biologic therapy/immune therapy/targeted therapy/other oncologic therapy regimen is adequately represented in the published evidence.
 - 3. Whether the reported study outcomes represent clinically meaningful outcomes experienced by patients. Generally, the definition of Clinically Meaningful outcomes are those recommended by ASCO, e.g., Hazard Ratio of less than 0.80 and the recommended survival benefit for OS and PFS should be at least 3 months.
 - 4. Whether the experimental design, in light of the drugs and conditions under investigation, is appropriate to address the investigative question. (For example, in some clinical studies, it may be unnecessary or not feasible to use randomization, double blind trials, placebos, or crossover).
 - 5. That non-randomized clinical trials with a significant number of subjects may be a basis for supportive clinical evidence for determining accepted uses of drugs.
 - 6. That case reports are generally considered uncontrolled and anecdotal information and do not provide adequate supportive clinical evidence for determining accepted uses of drugs.
 - 7. That abstracts (including meeting abstracts) without the full article from the approved peer-reviewed journals lack supporting clinical evidence for determining accepted uses of drugs.

IV. MEDICATION MANAGEMENT

- A. Please refer to the FDA label/package insert for details regarding these topics.

V. APPROVAL AUTHORITY

- A. Review – Utilization Management Department
- B. Final Approval – Utilization Management Committee

VI. ATTACHMENTS

- A. None

VII. REFERENCES

- A. Brock PR, et al. Sodium Thiosulfate for Protection from Cisplatin-Induced Hearing Loss. N Engl J Med. 2018 Jun 21;378(25):2376-2385.
- B. Freyer DR, et al. Effects of sodium thiosulfate versus observation on development of cisplatin-induced hearing loss in children with cancer (ACCL0431): a multicentre, randomised, controlled, open-label, phase 3 trial. Lancet Oncol. 2017 Jan;18(1):63-74.



EmblemHealth®

ConnectiCare

- C. Pedmark prescribing information. Fennec Pharmaceuticals Inc. Hoboken, NJ 2023.
- D. AHFS Drug Information. American Society of Health-Systems Pharmacists or Wolters Kluwer Lexi-Drugs Bethesda, MD 2023.
- E. Clinical Pharmacology Elsevier Gold Standard 2023.
- F. Micromedex® Healthcare Series: Thomson Micromedex, Greenwood Village, CO 2023.
- G. National Comprehensive Cancer Network. Cancer Guidelines and Drugs and Biologics Compendium 2023.
- H. Ellis LM, et al. American Society of Clinical Oncology perspective: Raising the bar for clinical trials by defining clinically meaningful outcomes. J Clin Oncol. 2014 Apr 20;32(12):1277-80.
- I. Medicare Benefit Policy Manual Chapter 15 Covered Medical and Other Health Services: <https://www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/Downloads/bp102c15.pdf>.
- J. NCQA UM 2023 Standards and Elements.