

# Medical Policy: Autologous Chondrocyte Implantation



| POLICY NUMBER | EFFECTIVE DATE | APPROVED BY                    |
|---------------|----------------|--------------------------------|
| MG.MM.SU.37   | 8/8/2025       | MPC (Medical Policy Committee) |

## ***IMPORTANT NOTE ABOUT THIS MEDICAL POLICY:***

Property of ConnectiCare, Inc. All rights reserved. The treating physician or primary care provider must submit to ConnectiCare, Inc. the clinical evidence that the patient meets the criteria for the treatment or surgical procedure. Without this documentation and information, ConnectiCare will not be able to properly review the request for prior authorization. This clinical policy is not intended to pre-empt the judgment of the reviewing medical director or dictate to health care providers how to practice medicine. Health care providers are expected to exercise their medical judgment in rendering appropriate care. The clinical review criteria expressed below reflects how ConnectiCare determines whether certain services or supplies are medically necessary. ConnectiCare established the clinical review criteria based upon a review of currently available clinical information (including clinical outcome studies in the peer-reviewed published medical literature, regulatory status of the technology, evidence-based guidelines of public health and health research agencies, evidence-based guidelines and positions of leading national health professional organizations, views of physicians practicing in relevant clinical areas, and other relevant factors). ConnectiCare, Inc. expressly reserves the right to revise these conclusions as clinical information changes, and welcomes further relevant information. Identification of selected brand names of devices, tests and procedures in a medical coverage policy is for reference only and is not an endorsement of any one device, test or procedure over another. Each benefit plan defines which services are covered. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that this service or supply is covered and/or paid for by ConnectiCare, as some plans exclude coverage for services or supplies that ConnectiCare considers medically necessary. If there is a discrepancy between this guideline and a member's benefits plan, the benefits plan will govern. In addition, coverage may be mandated by applicable legal requirements of the State of CT and/or the Federal Government. Coverage may also differ for our Medicare members based on any applicable Centers for Medicare & Medicaid Services (CMS) coverage statements including National Coverage Determinations (NCD), Local Coverage Determinations (LCD) and/or Local Medical Review Policies (LMRP). All coding and web site links are accurate at time of publication.

## **Definitions**

MACI® (autologous cultured chondrocytes on porcine collagen membrane) — an autologous cellularized scaffold product indicated for the repair of symptomatic single or multiple full-thickness cartilage defects of the knee with or without bone involvement in adults. MACI is a cellular sheet that consists of autologous chondrocytes seeded on a 3 x 5 cm, resorbable porcine Type I/III collagen membrane. The active ingredients of MACI are the autologous cultured chondrocytes and porcine Type I/III collagen.

## **Related Medical Guideline**

[Osteochondral Grafting](#)

## **Guideline**

Autologous chondrocyte implantation (ACI) is considered medically necessary for the repair of symptomatic, cartilaginous defects of the knee caused by acute or repetitive trauma when **all** of the following criteria are met:

1. Age 15–55 years of age
2. Presence of Grade III or Grade IV full thickness articular cartilage loss on a weight-bearing surface of the femoral condyle (medial or lateral femoral condyle)

3. Lesion is symptomatic (defined as lesion-related pain, swelling, or catching/locking that limits activities of daily living)
4. The focal chondral defect size is between 1–10 cm
5. Inadequate response to prior arthroscopic or other surgical repair procedure (e.g., debridement, microfracture, drilling/abrasion arthroplasty or osteochondral allograft/autograft)
6. The knee must be stable and aligned (a corrective procedure in combination with or prior to ACI may be necessary to ensure stability, alignment, and normal weight distribution within the joint)
7. Prior to the procedure, there must be an expectation that the member will be able to fully participate in a prescribed post-op rehabilitation program necessary to insure optimal outcome

**Note: Members can develop a new injury and another cartilage defect in the same knee. A subsequent or second procedure for a different defect of the same knee may be deemed medically necessary only if it is performed at least 6 months after the prior or initial procedure on that knee.**

## Documentation

All of the following must be documented and made available to the plan upon request:

1. Signs and symptoms limiting activities of daily living (pain, swelling, locking, crepitus, catching, giving-way, etc.)
2. Dates and outcomes of all previous surgical procedures on the knee (chondroplasty, drilling, microfracture, debridement, abrasion, etc.)
3. Pre-operative confirmation that the knee is stable and aligned with normal or optimal weight distribution within the joint
4. The location, dimensions, grade and depth of the identified focal chondral lesion, the condition of the surrounding articular cartilage

Documentation must support that prior to the procedure there was an expectation that the member would be able to fully participate in the post-op rehabilitation program necessary to insure optimal outcome.

## Limitations/Exclusions

1. ACI is not considered medically necessary for the following indications due to insufficient evidence of therapeutic value:
  - Kissing lesions (includes degeneration or disease on the femoral and tibial aspect of the joint)
  - History of patellar/multiple defects
  - In joints other than the knee
2. Repeat ACI for the same lesion is not considered medically necessary, as it has not been sufficiently studied
3. ACI is not considered reasonable and medically necessary in the following circumstances:

- As initial or first line surgical therapy
- In a member who has had a previous total meniscectomy (Note: presence of an unstable or torn meniscus requires partial resection, repair or replacement prior to or concurrent with MACI implantation. MACI is not recommended in patients with a total meniscectomy)
- In a member with a known history of anaphylaxis to gentamicin or sensitivities to materials of bovine origin
- In a member with infection at any of the proposed operative sites
- In a member with a cartilaginous defect associated with osteoarthritis or inflammatory diseases or where an osteoarthritic or inflammatory process significantly and adversely affects the quality of the peri-lesional cartilage
- > 55 years of age (Clinical trials of MACI did not include subjects over the age of 55)

## Procedure Codes

|       |   |
|-------|---|
| J7330 | Autologous cultured chondrocytes, implant   |
| S2112 | Arthroscopy, knee, surgical for harvesting of cartilage (chondrocyte cells) (Commercial only) |
| 27412 | Autologous chondrocyte implantation, knee   |

## Diagnosis Codes

|         |   |
|---------|---|
| M24.10  | Other articular cartilage disorders, unspecified site |
| M25.861 | Other specified joint disorders, right knee           |
| M25.862 | Other specified joint disorders, left knee            |
| M25.869 | Other specified joint disorders, unspecified knee     |
| M94.8X8 | Other specified disorders of cartilage, other site    |
| M94.9   | Disorder of cartilage, unspecified                    |

## References

Brittberg M, Lindahl A, Nilsson A, Ohlsson C, Isaksson O, Peterson L. Treatment of deep cartilage defects in the knee with autologous chondrocyte transplantation. *N Engl J Med*. 1994;331:889-895.

MACI [product insert]. Cambridge, Mass: Vericel Corporation; 2016.

Cole BJ, DiMasi M. Case report #10: single-stage autologous chondrocyte implantation and lateral meniscus allograft reconstruction. *Orthopedic Technology Review*. March 2000;2(3).

Gillogly SD, Voight M, Blackburn T. Treatment of articular cartilage defects of the knee with autologous chondrocyte implantation. *J Orthop Sports Phys Ther*. 1998;28(4):241-251.

Mandelbaum B, Browne J, Erggelet C, Fu F, Micheli L, Moseley JB. 4-year multicenter outcome of autologous chondrocyte implantation of the knee. Presented at: The Annual Meeting of the American Academy of Orthopedic Surgeons; March 2001; San Francisco, Calif.

Medical Policy:  
Autologous Chondrocyte Implantation



Mandelbaum BR, Browne JE, Fu F, et al. Articular cartilage lesions of the knee. *Am J Sports Med.* 1998;26:853-861.

Minas T. Chondrocyte implantation in the repair of chondral lesions of the knee: economics and quality of life. *Am J Orthop.* 1998;27:739-744.

Peterson L. Autologous chondrocyte transplantation: 2-10 year follow-up in 219 patients. Presented at: The Annual Meeting of the American Academy of Orthopedic Surgeons; March 1998; New Orleans, La.

Simonian PT, Sussman PS, Wickiewicz TL, Paletta GA, Warren RF. Contact pressures at osteochondral donor sites in the knee. *Am J Sports Med.* 1998;26:491-494.

Specialty-matched clinical peer review.

Wroble RR. Articular cartilage injury and autologous chondrocyte implantation: which patients might benefit? *The Physician and Sportsmedicine.* November 2000;28(11).

Revision History

| DATE          | REVISION  |
|---------------|---|
| Aug. 8, 2025  | Transferred policy content to individual company branded template   |
| Aug. 11, 2023 | Removed trochlear aspect, as a component of the femoral condyle, from the indication pertaining to articular cartilage loss   |
| Jan. 14, 2022 | ConnectiCare retired MCG criteria for this service and adopted the clinical criteria of its parent corporation EmblemHealth   |
| Aug. 14, 2020 | Removed "Patellofemoral disorders" from the list of conditions for which ACI is not considered medically necessary, as it is already excluded under "History of patellar/multiple defects." |
| Oct. 13, 2017 | Replaced Carticel (removed from the market) with MACI   |