

PHARMACY PRE-AUTHORIZATION CRITERIA

DRUG (S)	Xermelo (telotristat ethyl)
POLICY #	21106
INDICATIONS	Xermelo is indicated for the treatment of carcinoid syndrome diarrhea in combination with somatostatin analog (SSA) therapy in adults inadequately controlled by SSA therapy.
CRITERIA	<p>Xermelo is covered only if the following prior authorization criteria are met:</p> <ul style="list-style-type: none"> • The patient is 18 years or older <p>AND</p> <ul style="list-style-type: none"> • The patient has been on a long-acting somatostatin analog (SSA) therapy (e.g. Somatuline Depot [lanreotide for injection], Sandostatin LAR Depot [octreotide for injection], for at least 3 consecutive months <p>AND</p> <ul style="list-style-type: none"> • While on long-acting somatostatin analog therapy (prior to starting Xermelo), the patient continues to have at least four bowel movements per day <p>AND</p> <ul style="list-style-type: none"> • Xermelo will be used in combination with a long-acting somatostatin analog therapy <p>AND</p> <ul style="list-style-type: none"> • Prescribed by, or in consultation with, an oncologist or gastroenterologist for patients with a diagnosis of refractory carcinoid syndrome diarrhea
LIMITATIONS	<p>Xermelo will not be approved in treatment naïve patients. Xermelo will not be approved as monotherapy.</p> <p>If approved, an authorization will be granted for 12 weeks.</p> <p>For reauthorization, documentation showing a decrease in the number of bowel movements per day is required.</p>
REFERENCES	<ol style="list-style-type: none"> 1. Facts & Comparisons Online 2. Xermelo™ tablets [prescribing information]. The Woodlands, TX: Merck; February 2017.
P&T REVIEW HISTORY	5/17, 1/18
REVISION RECORD	1/18