

Medical Policy:

Cerezyme® (imiglucerase) intravenous infusion

POLICY NUMBER	LAST REVIEW	ORIGIN DATE
MG.MM.PH.223	March 7, 2025	2019

Medical Guideline Disclaimer Property of EmblemHealth. All rights reserved.

The treating physician or primary care provider must submit to EmblemHealth, or ConnectiCare, as applicable (hereinafter jointly referred to as "EmblemHealth"), the clinical evidence that the member meets the criteria for the treatment or surgical procedure. Without this documentation and information, EmblemHealth will not be able to properly review the request preauthorization or post-payment review. The clinical review criteria expressed below reflects how EmblemHealth determines whether certain services or supplies are medically necessary. This clinical policy is not intended to pre-empt the judgment of the reviewing medical director or dictate to health care providers how to practice medicine. Health care providers are expected to exercise their medical judgment in rendering appropriate care.

EmblemHealth established the clinical review criteria based upon a review of currently available clinical information (including clinical outcome studies in the peer reviewed published medical literature, regulatory status of the technology, evidence-based guidelines of public health and health research agencies, evidence-based guidelines and positions of leading national health professional organizations, views of physicians practicing in relevant clinical areas, and other relevant factors). EmblemHealth expressly reserves the right to revise these conclusions as clinical information changes and welcomes further relevant information. Each benefit program defines which services are covered. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that this service or supply is covered and/or paid for by EmblemHealth, as some programs exclude coverage for services or supplies that EmblemHealth considers medically necessary.

If there is a discrepancy between this guideline and a member's benefits program, the benefits program will govern. Identification of selected brand names of devices, tests and procedures in a medical coverage policy is for reference only and is not an endorsement of any one device, test or procedure over another. In addition, coverage may be mandated by applicable legal requirements of a state, the Federal Government or the Centers for Medicare & Medicaid Services (CMS) for Medicare and Medicaid members. All coding and web site links are accurate at time of publication.

EmblemHealth may also use tools developed by third parties, such as the MCG™ Care Guidelines, to assist us in administering health benefits. The MCG™ Care Guidelines are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice. EmblemHealth Services Company, LLC, has adopted this policy in providing management, administrative and other services to EmblemHealth Plan, Inc., EmblemHealth Insurance Company, EmblemHealth Services Company, LLC, and Health Insurance Plan of Greater New York (HIP) related to health benefit plans offered by these entities. ConnectiCare, an EmblemHealth company, has also adopted this policy. All of the aforementioned entities are affiliated companies under common control of EmblemHealth Inc.

Definitions

Cerezyme, an analogue of β -glucocerebrosidase, is indicated for the long-term enzyme replacement therapy for patients with a confirmed diagnosis of Type 1 Gaucher disease that results in at least one of the following: anemia, thrombocytopenia, bone disease, hepatomegaly, or splenomegaly.

Length of Authorization

Coverage will be provided for 12 months and may be renewed.

Dosing Limits [Medical Benefit]

Each individual dose must not exceed 60 U/kg administered intravenously no more frequently than three times per week.

Max Units (per dose and over time) [HCPCS Unit]:

720 billable units every 14 days

Guideline

I. INITIAL CRITERIA

1. Type 1 Gaucher Disease

Approve if the patient meets the following criteria:

- A. Patient is at least 2 years of age; AND
- B. Used as a single agent; AND
- C. Patient has a documented diagnosis of Type 1 Gaucher Disease confirmed by one of the following:
 - i. Significantly reduced or absent glucocerebrosidase enzyme activity as measured by a betaglucosidase leukocyte (BGL) test
 - ii. Detection of mutations in the glucocerebrosidase (GBA) gene; ANDPatient has Type 1 Gaucher disease; **AND**
- D. Patient's disease results in one or more of the following
 - i. Anemia-related symptoms [i.e., blood transfusion dependency and/or hemoglobin ≤ 11 g/dL (women and children) or ≤ 12 g/dL (men)] **OR**
 - ii. Thrombocytopenia (platelet count ≤ 120,000/mm3) **OR**
 - iii. Hepatomegaly or splenomegaly OR
 - iv. Skeletal disease **OR** (e.g., lesions, remodeling defects and/or deformity of long bones, osteopenia/osteoporosis, etc.)
 - v. Symptomatic disease (e.g., bone pain, fatigue, dyspnea, abdominal distension, diminished quality of life, etc.) **AND**
- E. Cerezyme is prescribed by or in consultation with a geneticist, endocrinologist, a metabolic disorder subspecialist, or a physician who specializes in the treatment of lysosomal storage disorders.

II. RENEWAL CRITERIA

- 1. Patient continues to meet Initial criteria; AND
- 2. Disease response with treatment; AND
- 3. Absence of unacceptable toxicity from the drug. *Examples of unacceptable toxicity include: hypersensitivity reactions, etc.*

Applicable Procedure Codes

Code	Description	
J1786	Injection, imiglucerase, 10 units	

Applicable NDCs

Code	Description
58468-4663-01	Cerezyme 400-unit Solution Reconstituted

ICD-10 Diagnoses

Code	Description
E75.22	Gaucher Disease

Revision History

Company(ies)	DATE	REVISION
--------------	------	----------

EmblemHealth & ConnectiCare	3/7/2025	Annual Update: Update initial criteria guidelines with specific indications. Removed The diagnosis is established by one of the following (i or ii): i. Demonstration of deficient β-glucocerebrosidase activity in leukocytes or fibroblasts; OR ii. Molecular genetic testing documenting glucocerebrosidase gene mutation; AND. Addition of at least 2 years of age and used as a single agent. Update billable units from 700 to 720
EmblemHealth & ConnectiCare	3/25/2024	Annual Review: Updated dosing limits, added renewal criteria, updated ICD from E75.2 to E75.22
EmblemHealth & ConnectiCare	4/10/2023	Transfer from CCUM template to medical CoBranded template Retired MG.MM.PH.73
EmblemHealth & ConnectiCare	3/16/2022	Annual Revision: no criteria changes
EmblemHealth & ConnectiCare	3/17/2021	Annual Revision: Gaucher Disease: Moved the designation of "Type 1" disease from the indication to criteria

References

1. Cerezyme® intravenous infusion [prescribing information]. Cambridge, MA: Genzyme; July 2024