

Other Uses with Supportive Evidence

- 2. Pervasive Developmental Disorders (e.g., autism spectrum disorder, Asperger's disorder).** Approve for 1 year if the patient has symptoms of attention deficit hyperactivity disorder (e.g., inattention, hyperactivity).

II. Coverage of Qelbree is recommended in those who meet the following criteria:

FDA-Approved Indication

- 1. Attention Deficit Hyperactivity Disorder.** Approve for 1 year if the patient is ≥ 6 years of age.

CONDITIONS NOT RECOMMENDED FOR APPROVAL

Coverage of atomoxetine capsules (Strattera, generic), clonidine ER tablets (Kapvay, generic), guanfacine ER tablets (Intuniv, generic), or Qelbree is not recommended in the following situations:

- 1. Binge-Eating Disorder.** In one 10-week, placebo-controlled study in outpatients with binge-eating disorder (n = 40), atomoxetine was associated with a significantly greater reduction in binge-eating episode frequency vs. placebo.⁷ Additional studies with atomoxetine are needed. There are no data with guanfacine ER tablets, clonidine ER tablets, or Qelbree.
- 2. Depression without Attention Deficit/Hyperactivity Disorder.** Limited information is available on the use of atomoxetine for the treatment of major depressive disorder. In three case reports and one case series in 15 patients with depressive disorders, adding atomoxetine to a selective serotonin reuptake inhibitor resulted in further improvement.^{8,9} However, in a published controlled trial, patients with major depressive disorder (without ADHD) [n = 276] were treated with sertraline at doses up to 200 mg/day.¹⁰ Patients who continued to experience depressive symptoms (n = 146) were then randomly assigned to either treatment with atomoxetine 40 to 120 mg/day or placebo for an additional 8 weeks. There was no difference between the atomoxetine/sertraline and placebo/sertraline treatment groups in mean change in depressive symptom severity or in the number of patients whose depressive symptoms remitted (40.3% vs. 37.8%, respectively; P = 0.865). Atomoxetine did not improve clinically significant depression in patients with Parkinson disease (n = 55) in one study.¹¹ There are no data with guanfacine ER tablets, clonidine ER tablets, or Qelbree.
- 3. Fibromyalgia.** In case reports, atomoxetine was effective in reducing fatigue and pain in fibromyalgia syndrome.¹² Well-controlled trials with atomoxetine are needed to establish safety and efficacy. There are no data with guanfacine ER tablets, clonidine ER tablets, or Qelbree.
- 4. Improve Cognitive Function (or Neuroenhancement).** The use of prescription medication to augment cognitive or affective function in otherwise healthy individuals (also known as neuroenhancement) is increasing in adult and pediatric populations.²⁰ A 2013 Ethics, Law, and Humanities Committee position paper, endorsed by the American Academy of Neurology, indicates that based on currently available data and the balance of ethics issues, neuroenhancement in children and adolescents without a diagnosis of a neurologic disorder is not justifiable. The prescription of neuroenhancements is inadvisable due to numerous social, developmental, and professional integrity issues. Several studies have evaluated atomoxetine for cognitive function in various patient populations, including patients with Huntington disease¹², Alzheimer's disease¹⁴, schizophrenia^{15,16}, and Parkinson's disease.¹⁷ However, atomoxetine has not demonstrated clinical benefit.

5. **Long-Term Combination Therapy (i.e., > 2 months) with atomoxetine (Strattera, generic) and Central Nervous System (CNS) Stimulants used for the Treatment of Attention Deficit/Hyperactivity Disorder (e.g., mixed amphetamine salts ER capsules [Adderall XR, generic], methylphenidate ER tablets, methylphenidate immediate-release tablets).** Currently, data do not support using atomoxetine and CNS stimulant medications concomitantly.^{18,19} Short-term drug therapy (2 months or less) with both atomoxetine and CNS stimulant medications is allowed for transitioning the patient to only one drug. Guanfacine ER tablets and clonidine ER tablets are indicated for use as monotherapy or as adjunctive therapy to CNS stimulant medications; therefore, long-term combination therapy with either agent and CNS stimulants is appropriate.²⁻³ Qelbree labeling does not address combination use with CNS stimulants at this time.⁴
6. **Nocturnal Enuresis.** In case reports, children with ADHD and other comorbid psychiatric diagnoses who had nocturnal enuresis and were treated with atomoxetine had resolution of their enuresis.²¹ In one controlled trial in pediatric patients (n = 87) with nocturnal enuresis, atomoxetine increased the average number of dry nights per week by 1.47 vs. 0.60 for placebo (P = 0.01).²² Additional controlled trials with atomoxetine are needed. Data with guanfacine ER tablets, clonidine ER tablets, or Qelbree are lacking.
7. **Weight Loss.** In one 12-week, placebo-controlled study in obese women (n = 30), atomoxetine resulted in a mean -3.7% loss vs. 0.2% gain with placebo when combined with a hypocaloric diet (500 kcal/day deficit).²³ Atomoxetine did not demonstrate efficacy for weight reduction in patients with schizophrenia (n = 37) treated with antipsychotics (clozapine or olanzapine).²⁴ Additional studies are needed.
8. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

REFERENCES

1. Strattera® capsules [prescribing information]. Indianapolis, IN: Lilly; January 2022.
2. Intuniv® extended-release tablets [prescribing information]. Lexington, MA: Shire; December 2019.
3. Kapvay® extended-release tablets, oral [prescribing information]. Overland Park, KS: Concordia; February 2020.
4. Qelbree® extended-release capsules [prescribing information]. Rockville, MD: Supernus; April 2022.
5. Nash K, Carter KJ. Treatment options for the management of pervasive developmental disorders. *Int J Psychiatry Med.* 2016;51(2):201-210.
6. Goel R, Hong JS, Findling RL. An update on pharmacotherapy of autism spectrum disorder in children and adolescents. *Int Rev Psychiatry.* 2018;30(1):78-95.
7. McElroy SL, Guerdjikova A, Kotwal R, et al. Atomoxetine in the treatment of binge-eating disorder: a randomized placebo-controlled trial. *J Clin Psychiatry.* 2007;68:390-398.
8. Berigan TR. Atomoxetine used adjunctively with selective serotonin reuptake inhibitors to treat depression. *Prim Care Companion J Clin Psychiatry.* 2004;6:93-94.
9. Carpenter LL, Milosavljevic N, Jordan JD, Schechter JM, Tyrka AR, Price LH. Augmentation with open-label atomoxetine for partial or nonresponse to antidepressants. *J Clin Psychiatry.* 2005;66:1234-1238.
10. Michelson D, Adler LA, Amsterdam JD, et al. Addition of atomoxetine for depression incompletely responsive to sertraline: a randomized, double-blind, placebo-controlled study. *J Clin Psychiatry.* 2007;68:582-587.
11. Weintraub D, Mavandadi S, Mamikonyan E, et al. Atomoxetine for depression and other neuropsychiatric symptoms in Parkinson disease. *Neurology.* 2010;75:448-455.
12. Berigan T. The use of atomoxetine adjunctively in fibromyalgia syndrome. *Can J Psychiatry.* 2004;49:499-500.
13. Beglinger LJ, Adams WH, Paulson H, et al. Randomized controlled trial of atomoxetine for cognitive dysfunction in early Huntington disease. *J Clin Psychopharmacol.* 2009;29:484-487.
14. Mohs R, Shiovitz TM, Tariot PN, et al. Atomoxetine augmentation of cholinesterase inhibitor therapy in patients with Alzheimer disease: 6-month, randomized, double-blind, placebo-controlled, parallel-trial study. *Am J Geriatr Psychiatry.* 2009;17:752-759.
15. Kelly DL, Buchanan RW, Boggs DL, et al. A randomized double-blind trial of atomoxetine for cognitive impairments in 32 people with schizophrenia. *J Clin Psychiatry.* 2009;70:518-525.



16. Friedman JI, Carpenter D, Lu J, et al. A pilot study of adjunctive atomoxetine treatment to second-generation antipsychotics for cognitive impairment in schizophrenia. *J Clin Psychopharmacol*. 2008;28:59-63.
17. Marsh L, Biglan K, Gerstenhaber M, Williams JR. Atomoxetine for the treatment of executive dysfunction in Parkinson's disease: a pilot open-label study. *Mov Disord*. 2009;30:277-282.
18. Treuer T, Gau SS-F, Mendez L, et al. A systematic review of combination therapy with stimulants and atomoxetine for attention-deficit/hyperactivity disorder, including patient characteristics, treatment strategies, effectiveness, and tolerability. *J Child Adolesc Psychopharmacol*. 2013;23(3):179-193.
19. Clemow DB, Mason OW, Sarkis EH, et al. Atomoxetine monotherapy compared with combination therapy for the treatment of ADHD: a retrospective chart review study. *Expert Rev Neurother*. 2015;15(11):1353-1366.
20. Graf WD, Nagel SK, Epstein LG, et al. Pediatric neuroenhancement: ethical, legal, social, and neurodevelopmental implications. *Neurology*. 2013;80:1251-1260.
21. Shatkin JP. Atomoxetine for the treatment of pediatric nocturnal enuresis. *J Child Adolesc Psychopharmacol*. 2004;14(3):443-447.
22. Sumner CR, Schuh KJ, Sutton VK, et al. Placebo-controlled study of the effects of atomoxetine on bladder control in children with nocturnal enuresis. *J Child Adolesc Psychopharmacol*. 2006;16:699-711.
23. Gadde KM, Yonish GM, Wagner HR, et al. Atomoxetine for weight reduction in obese women: a preliminary randomized controlled trial. *Int J Obes (Lond)*. 2006;30:1138-1142.
24. Ball MP, Warren KR, Feldman S, et al. Placebo-controlled trial of atomoxetine for weight reduction in people with schizophrenia treated with clozapine or olanzapine. *Clin Schizophr Relat Psychoses*. 2011;5:17-25.