

## PRIOR AUTHORIZATION POLICY

**POLICY:** Cardiology – Corlanor Prior Authorization Policy

- Corlanor® (ivabradine tablets and oral solution – Amgen)

**REVIEW DATE:** 06/29/2022

---

### OVERVIEW

Corlanor, a hyperpolarization-activated cyclic nucleotide-gated channel blocker, is indicated for the following uses:<sup>1</sup>

- **Heart failure, in adults**, to reduce the risk of hospitalization for worsening of the disease in those with stable, symptomatic chronic heart failure with left ventricular ejection fraction (LVEF)  $\leq 35\%$ , who are in sinus rhythm with a resting heart rate  $\geq 70$  beats per minute (bpm) and either are receiving maximally tolerated doses of beta blockers or have a contraindication to beta blocker use.
- **Heart failure, in pediatric patients  $\geq 6$  months and older**, for treatment of stable symptomatic disease due to dilated cardiomyopathy, among those who are in sinus rhythm with an elevated heart rate.

Data are available with Corlanor that note improvement in symptoms and increased exercise performance in patients with inappropriate sinus tachycardia, defined as a sinus heart rate  $> 100$  bpm at rest (with a mean 24-hour heart rate  $> 90$  bpm not due to primary causes) which is generally associated with distressing symptoms such as palpitations, weakness, dizziness and syncope.<sup>2-9</sup> Beta blockers have also been used for this condition. Limited data are available for other treatments that have been used and/or effectiveness have not been established (e.g., beta blockers, fludrocortisone, volume expansion, clonidine, and erythropoietin).

### Guidelines

A few guidelines have recommendations that involve Corlanor.

- **Heart Failure:** In 2022 the American Heart Association/American College of Cardiology/Heart Failure Society of America published guidelines in 2022 for the management of heart failure.<sup>10</sup> For patients with symptomatic (New York Heart Association Class II to III) stable chronic heart failure with reduced ejection fraction (LVEF  $\leq 35\%$ ) who are receiving guideline-directed medical therapy, including a beta blocker at maximum tolerated dose, and who are in sinus rhythm with a heart rate of  $\geq 70$  beats per minute at rest, Corlanor can be beneficial to reduce heart failure hospitalizations and cardiovascular death.
- **Inappropriate Sinus Tachycardia:** The 2015 Heart Rhythm Society Expert Consensus Statement on the diagnosis and treatment of postural tachycardia syndrome, inappropriate sinus tachycardia, and vasovagal syncope state that Corlanor can be useful for treating patients with inappropriate sinus tachycardia.<sup>2</sup> Additionally, the 2015 American College of Cardiology, American Heart Association Task Force on Clinical Practice Guidelines, and the Heart Rhythm Society also state that Corlanor is reasonable for ongoing management in patients with symptomatic inappropriate sinus tachycardia (class IIa recommendation).<sup>3</sup> Beta blockers may be considered for ongoing management in patients with symptomatic inappropriate sinus tachycardia (class IIb recommendation). Also, the guidelines state that the combination of beta blockers and Corlanor may be considered for the ongoing management of patients with inappropriate sinus tachycardia (class IIb recommendation). Because of the specialized skills required for evaluation and diagnosis of patients treated with Corlanor as well as the monitoring required for adverse events and long-term efficacy, approval requires Corlanor to be prescribed by or in consultation with a physician who specializes in the condition being treated.

### **POLICY STATEMENT**

Prior Authorization is recommended for prescription benefit coverage of Corlanor. All approvals are provided for the duration noted below.

**Automation:** None.

### **RECOMMENDED AUTHORIZATION CRITERIA**

Coverage of Corlanor is recommended in those who meet one of the following criteria:

#### **FDA-Approved Indications**

- 1. Heart Failure.** Approve for 1 year if the patient meets the following criteria (A, B, C, D, and E):
  - A) Patient is  $\geq 18$  years of age; AND
  - B) Patient has a left ventricular ejection fraction (LVEF)  $\leq 35\%$  currently or prior to initiation of Corlanor therapy; AND
  - C) Patient is in normal sinus rhythm or sinus tachycardia with a resting heart rate of  $\geq 70$  beats per minute; AND
  - D) Patient meets one of the following (i or ii):
    - i. Patient has tried or is currently receiving one beta blocker or heart failure treatment; OR  
Note: Examples of beta blockers are metoprolol succinate sustained-release, carvedilol, bisoprolol, and Coreg CR (carvedilol extended-release capsules).
    - ii. Patient has a contraindication to use of beta blocker therapy; AND  
Note: Examples that are contraindications to use of beta blockers are bronchospastic disease such as chronic obstructive pulmonary disease and asthma, severe hypotension or bradycardia.
  - E) Medication is prescribed by, or in consultation with, a cardiologist.
- 2. Heart Failure due to Dilated Cardiomyopathy in Pediatric Patients.** Approve for 1 year if the patient meets the following (A and B):
  - A) Patient is  $< 18$  years of age; AND
  - B) Medication is prescribed by, or in consultation with, a cardiologist.

#### **Other Uses with Supportive Evidence**

- 3. Inappropriate Sinus Tachycardia.** Approve for 1 year if the medication is prescribed by, or in consultation with, a cardiologist.

### **CONDITIONS NOT RECOMMENDED FOR APPROVAL**

Coverage of Corlanor is recommended in those who meet the following criteria:

- 1. Stable Angina Pectoris, in Patients Without Chronic Heart Failure.** Corlanor has been studied as a treatment for stable angina pectoris but further data are needed.<sup>11-13</sup> US guidelines addressing stable angina do not include Corlanor.<sup>14,15</sup>
- 2.** Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.



EmblemHealth®

ConnectiCare

**REFERENCES**

1. Corlanor® tablets and oral solution [prescribing information]. Thousand Oaks: CA; Amgen; August 2021.
2. Sheldon RS, Grubb BP, Olshansky B, et al. 2015 Heart Rhythm Society Expert Consensus statement on the diagnosis and treatment of postural tachycardia syndrome, inappropriate sinus tachycardia, and vasovagal syncope. *Heart Rhythm*. 2015;12(6):e41-e63.
3. Page RL, Caldwell MA, Calkins H, et al. 2015 ACC/AHA/HRS guideline for the management of adult patients with supraventricular tachycardia. A report of the American College of Cardiology/American Heart Association Task Force on clinical practice guidelines, and the Heart Rhythm Society. *Circulation*. 2016;133:e506-e574. Available at: <http://circ.ahajournals.org/content/133/14/e506.full.pdf+html>. Accessed on June 22, 2022.
4. Abed HS, Fulcher J, Kilborn NJ, Keech AC. Inappropriate sinus tachycardia: focus on ivabradine. *Intern Med J*. 2016;46(8):875-883.
5. Oliphant CS, Owens RE, Bolorunduro OB, Jha SK. Ivabradine: a review of labeled and off-label uses. *Am J Cardiovasc Drugs*. 2016;16:337-347.
6. Ruzieh M, Moustafa A, Sabbagh E, et al. Challenges in treatment of inappropriate sinus tachycardia. *Curr Cardiol Rev*. 2018;14:42-44.
7. Mathew ST, Po SS, Thadani U. Inappropriate sinus tachycardia – symptoms and heart rate reduction with ivabradine: a pooled analysis of prospective studies. *Heart Rhythm*. 2018;15:240-247.
8. Koruth JS, Lala A, Pinne S, et al. The clinical use of ivabradine. *J Am Coll Cardiol*. 2017;70(14):1777-1784.
9. Bonnet D, Berger F, Jokinen E, et al. Ivabradine in children with dilated cardiomyopathy and symptomatic chronic heart failure. *J Am Coll Cardiol*. 2017;70:1262-1272.
10. Heidenreich PA, Bozkurt B, Aguilar D, et al. 2022 AHA/ACC/HFSA guideline for the management of heart failure: a report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation*. 2022;145(18):e895-e1032.
11. Werdan K, Ebelt H, Nuding S, et al. Ivabradine in combination with beta-blocker improves symptoms and quality of life in patients with stable angina pectoris: results from the ADDITIONS study. *Clin Res Cardiol*. 2012;101(5):365-373.
12. Perings S, Stockl G, Kelm M, et al, on behalf of the RESPONSIVE study investigators. Effectiveness and tolerability of ivabradine with or without concomitant beta-blocker therapy in patients with chronic stable angina in routine clinical practice. *Adv Ther*. 2016;33:1550-1564.
13. Glezer M, Vasyuk Y, Karpov Y. Efficacy of ivabradine in combination with beta-blockers versus uptitration of beta-blockers in patients with stable angina (CONTROL-2 Study). *Adv Ther*. 2018;35:341-352.
14. Amsterdam EA, Wenger NK, Brindis RG, et al. 2014 AHA/ACC guideline for the management of patients with non-ST-elevation acute coronary syndromes: executive summary. A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation*. 2014;30:2354-2394. Available at: <http://circ.ahajournals.org/content/130/25/2354>. Accessed on June 22, 2022.
15. Fihn SD, Gardin JM, Abrams J, et al. 2012 ACCF/AHA/AATS/PCNA/SCAI/STS guideline for the diagnosis and management of patients with stable ischemic heart disease: executive summary. A report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines, and the American College of Physicians, American Association for Thoracic Surgery, Preventive Cardiovascular Nurses Association, Society for Cardiovascular Angiography and Interventions, and the Society of Thoracic Surgeons. *Circulation*. 2012;126:3097-3137. Available at: <http://circ.ahajournals.org/content/126/25/3097.full.pdf+html>. Accessed on June 22, 2022.